



Remarks on the nomenclature of the genera *Ptychodus* Agassiz, 1834 and *Buffonites* Sternberg, 1829 (Ptychodontidae, Chondrichthyes)

Luca GIUSBERTI, Manuel AMADORI, Jacopo AMALFITANO, Giorgio CARNEVALE & Jürgen KRIWET

L. Giusberti, Dipartimento di Geoscienze, Università di Padova, Via Gradenigo 6, I-35131 Padova, Italy; luca.giusberti@unipd.it
 M. Amadori, Via Pilone 91/A, I-61030 Lucrezia di Cartoceto (PU), Italy; manu.ama89@gmail.com
 J. Amalfitano, Dipartimento di Geoscienze, Università di Padova, Via Gradenigo 6, I-35131 Padova, Italy; jacopo.amalfitano@phd.unipd.it
 G. Carnevale, Dipartimento di Scienze della Terra, Università di Torino, Via Valperga Caluso 35, I-10125 Torino, Italy; giorgio.carnevale@unito.it
 J. Kriwet, Department of Palaeontology, Geozentrum, University of Vienna, Althanstraße 14, 1090 Vienna, Austria; juergen.kriwet@univie.ac.at

SUPPLEMENTARY ONLINE MATERIAL

LIST OF PUBLICATIONS IN THE LAST 50 YEARS (1968-2018) IN WHICH THE GENUS PTYCHODUS AGASSIZ, 1834 HAS BEEN USED AS A VALID TAXON

- Adnet S., Cappetta H. & Mertiniene R. (2008). Re-evaluation of squaloid shark records from the Albian and Cenomanian of Lithuania. *Cretaceous Research*, 29: 711-722.
- Albright L.B., Gillette D.D. & Titus A.L. (2013). Fossil vertebrates from the Tropic Shale (Upper Cretaceous), Southern Utah. In Titus A.L. & Loewen M.A. (eds), *At the top of the Grand Staircase: the Late Cretaceous of Southern Utah*, Indiana University Press, Bloomington and Indianapolis: 536-562.
- Antunes M.T. & Cappetta H. (2002). Sélaciens du Crétacé (Albien–Maastrichtien) d'Angola. *Palaeontographica*, A, 264: 85-146.
- Applegate S.P. (1970). The vertebrate fauna of the Selma Formation of Alabama, Part VIII. The Fishes. *Fieldiana: Geology Memoirs*, 8, 3: 383-433.
- Becker M.A., Wellner R.W., Mallery C.S. Jr. & Chamberlain J.A. Jr. (2010). Chondrichthyans from the Lower Ferron Sandstone Member of the Mancos Shale (upper Cretaceous: Middle Turonian) of Emery and Carbon Counties, Utah, USA. *Journal of Paleontology*, 84: 248-266.
- Bendix-Almgreen S.E. (1983). *Carcharodon megalodon* from the Upper Miocene of Denmark, with comments on elasmobranch tooth enameloid: coronoin. *Bulletin of the Geological Society of Denmark*, 32: 1-32.
- Blanco-Piñón A., Garibay-Romero L.M. & Alvarado-Ortega J. (2007). The oldest stratigraphic record of the Late Cretaceous shark *Ptychodus mortoni* Agassiz, from Vallecillo, Nuevo León, northeastern Mexico. *Revista Mexicana de Ciencias Geológicas*, 24: 25-30.
- Brignon A. (2015). Senior synonyms of *Ptychodus latissimus* Agassiz, 1835 and *Ptychodus mammillaris* Agassiz, 1835 (Elasmobranchii) based on teeth from the Bohemian Cretaceous Basin (the Czech Republic). *Acta Musei Nationalis Pragae, Series B-Historia Naturalis*, 71: 5-14.
- Brito P.M. & Janvier P. (2002). A ptychodontid (Chondrichthyes, Elasmobranchii) from the Upper Cretaceous of South America. *Geodiversitas*, 24: 785-790.
- Cappetta H. (1987). *Handbook of Paleichthyology*, Vol. 3B: Chondrichthyes II. 193 pp. Gustav Fischer Verlag, Stuttgart.
- Cappetta H. (ed.) (2012). Chondrichthyes. Mesozoic and Cenozoic Elasmobranchii: teeth. *Handbook of Paleichthyology*, volume 3E. 512 pp. Verlag Dr. Friedrich Pfeil, München.
- Cappetta H. & Case G.R. (1999). Additions aux faunes de sélaciens du Crétacé du Texas (Albien supérieur-Campanien). *Palaeo Ichthyologica*, 9: 5-111.
- Carrillo-Briceño J.D. & Lucas S.G. (2013). The first tooth set of *Ptychodus atcoensis* (Elasmobranchii: Ptychodontidae), from the Cretaceous of Venezuela. *Swiss Journal of Palaeontology*, 132: 69-75.
- Cigala Fulgosi F., Kotsakis T., Massari F., Medizza F. & Sorbini L. (1980). In I vertebrati fossili italiani, Catalogo della Mostra. Il giacimento di S. Anna d'Alfaedo: 123-129.
- Cuny G. (2008). Mesozoic hybodont sharks from Asia and their relationships to the genus *Ptychodus*. *Acta Geologica Polonica*, 58: 211-216.
- Dalla Vecchia F. M., Barbera C., Bizzarrini F., Bravi S., Delfino M., Giusberti L., Guidotti G., Mietto P., Papazzoni C.A., Roghi G., Signore M. & Simone O. (2005). Il Cretaceo marino. In Bonfiglio L. (ed.), *Palaeontologia dei Vertebrati in Italia. Memorie Museo Civico di Storia Naturale di Verona, Sezione Scienze della Terra*, 6: 101-116.
- Diedrich C.G. (2013). Facies related phylostratigraphy of the benthic neoselachian *Ptychodus* from the Late Cretaceous (Cenomanian/Turonian) of the Pre-North Sea Basin of Europe. *Cretaceous Research*, 41: 17-30.
- Ekrť B., Košťák M., Mazuch M., Voigt S. & Wiese F. (2004). New records of teleosts from the Late Turonian (Late Cretaceous) of the Bohemian Cretaceous Basin (Czech Republic). *Cretaceous Research*, 29: 659-673.
- Everhart M.J. (2005). *Oceans of Kansas: a natural history of the Western Interior Sea*. 322 pp. Indiana University Press, Bloomington and Indianapolis.
- Everhart M.J. & Caggiano T. (2004). An associated dentition and calcified vertebral centra of the Late Cretaceous elasmobranch, *Ptychodus anonymus* Williston 1900. *Paludicola*, 4: 125-136.
- Everhart M.J. & Darnell M.K. (2004). Occurrence of *Ptychodus mammillaris* (Elasmobranchii) in the Fairport Chalk Member of the Carlile Shale (Upper Cretaceous) of Ellis County, Kansas. *Transactions of the Kansas Academy of Science*, 107: 126-130.
- Fischer J., Kogan I., Popov E., Janetschke N. & Licht M. (2017). The Late Cretaceous chondrichthyan fauna of the Elbtal Group (Saxony, Germany). *Research & Knowledge*, 3: 13-17.

- Gagnaison C. (2013). Les assemblages de vertébrés dans deux sites paléontologiques du bassin miocène de Savigné-sur-Lathan/Noyant-sous-le-Lude: la Guimardière et Pelmer (Maine-et-Loire, France). *Geodiversitas*, 35: 67-103.
- Goto M., Uyeno T. & Yabumoto Y. (1996). Summary of Mesozoic elasmobranch remains from Japan. In Arratia G. & Viohl G. (eds), Mesozoic fishes – Systematics and Paleoeology. Verlag Dr. Friedrich Pfeil: 73-82.
- Guinot G., Underwood C.J., Cappetta H. & Ward D.J. (2013). Sharks (Elasmobranchii: Euselachii) from the Late Cretaceous of France and the UK. *Journal of Systematic Palaeontology*, 11: 589-671.
- Hakenberg M. (1969). Albian and Cenomanian between Małogoszcz and Staniewice, SW border of the Holy Cross Mountains. *Studia Geologica Polonica*, 26: 1-126.
- Hamm S. (2017). First Associated Tooth Set of *Ptychodus mammillaris* in North America, Pfeifer Shale Member (Lower Middle Turonian), Greenhorn Limestone. *Transactions of the Kansas Academy of Science*, 120: 17-30.
- Hamm S. & Cicimurri D.J. (2011). Early Coniacian (Late Cretaceous) Selachian fauna from the Basal Atco Formation, Lower Austin Group, North Central Texas. *Paludicola*, 8: 107-127.
- Herman J. (1977). Les sélections des terrains néocrétacés et paléocènes de Belgique et des contrées limitrophes. Elements d'une biostratigraphie inter-continentale. *Mémoires pour servir à l'Explication des Cartes géologiques et minières de la Belgique*, 15 [1975]: 1-401.
- Hoffman B.L., Hageman S.A. & Claycomb G.D. (2016). Scanning electron microscope examination of the dental enameloid of the Cretaceous durophagous shark *Ptychodus* supports neoselachian classification. *Journal of Paleontology*, 90: 741-762.
- Itoigawa J., Nishimoto N. & Hiroyuki A. (1977). Cretaceous fossil elasmobranchs from Japan (first report). *Bulletin of the Mizunami Fossil Museum*, 4: 119-138.
- Kauffman E.G. (1972). *Ptychodus* predation upon a Cretaceous *Inoceramus*. *Palaeontology*, 15: 439-444.
- Ladwig J. (2003). Erstnachweis eines Zahnes von *Ptychodus latissimus* Agassiz 1843 aus dem Geschiebe. *Der Geschiebsammler*, 36: 43-46.
- Longbottom A.E. & Patterson C. (1987). Fishes. In Smith A.B. (ed.), Fossils of the Chalk (Field Guides to Fossils, nr. 2), The Palaeontological Association: 238-265.
- Lucas S. (2006). Type locality of the Late Cretaceous shark *Ptychodus whipplei* Marcou, Northern New Mexico. *New Mexico Museum of Natural History and Science Bulletin*, 35: 157-159.
- MacLeod N. (1982). The first North American occurrence of the Late Cretaceous elasmobranch *Ptychodus rugosus* Dixon with comments on the functional morphology of the dentition and dermal denticles. *Journal of Paleontology*, 56: 403-409.
- Małeck J. (1980). Teeth of fishes of the genus *Ptychodus* from Cretaceous chalk sediments in the vicinity of Cracow. *Bulletin de l'Académie Polonaise des Sciences, Série des Sciences de la Terre*, 28: 51-57.
- Malvesy T., Cappetta H., Dutheil D.B., Otero O. & Oudoire T. (2002). Les collections du Musée d'Histoire Naturelle de Lille. IV. – poissons mésozoïques et cénozoïques (types et figurés). *Annales de la Société Géologique du Nord, série 2*, 9: 147-157.
- Marcinowski R. & Radwanski A. (1983). The Mid-Cretaceous transgression onto the Central Polish Uplands (marginal part of the Central European Basin). *Zitteliana*, 10: 65-95.
- McIntosh A.P., Shimada K. & Everhart M.J. (2016). Late Cretaceous marine vertebrate fauna from the Fairport Chalk Member of the Carlile Shale in southern Ellis County, Kansas, USA. *Transactions of the Kansas Academy of Science*, 119: 222-230.
- Mendiola C. (2004). Primera cita espa-ola del género *Ptychodus* Agassiz 1839 (Chondrichthyes, Euselachii). *Revista de la Societat Paleontològica d'Elx*, 13: 1-11.
- Mutter R.J., Iturralde-Vinent M. & Carmona J.F. (2005). The first Mesozoic Caribbean shark is from the Turonian of Cuba: *Ptychodus cyclodontis* sp. nov. (?Neoselachii). *Journal of Vertebrate Paleontology*, 25: 976-978.
- Niedźwiedzki R. & Kalina M. (2003). Late Cretaceous sharks in the Opole Silesia region (SW Poland). *Geologia Sudetica*, 35: 13-24.
- Popov E.V. & Lapkin A.V. (2000). A new shark species of the genus *Galeorhinus* (Chondrichthyes, Triakidae) from the Cenomanian of the Lower Volga River Basin. *Paleontological Journal*, 34: 435-438.
- Pyrah B.J. (1979). Catalogue of type and figured fossils in the Yorkshire Museum: Part 4. Pisces, Reptilia, Aves, Mammalia, Plantae. *Proceedings of the Yorkshire Geological Society*, 42: 415-437.
- Radwanski A. & Marcinowski R. (1996). Elasmobranch teeth from the mid-Cretaceous sequence of the Mangyshlak Mountains, Western Kazakhstan. *Acta Geologica Polonica*, 46: 165-169.
- Shimada K. (2012). Dentition of Late Cretaceous shark, *Ptychodus mortoni* (Elasmobranchii, Ptychodontidae). *Journal of Vertebrate Paleontology*, 32: 1271-1284.
- Shimada K. & Everhart M.J. (2003). *Ptychodus mammillaris* (Elasmobranchii) and *Enchodus* cf. *E. shumardi* (Teleostei) from the Fort Hays Limestone Member of the Niobrara Chalk (Upper Cretaceous) in Ellis County, Kansas. *Transactions of the Kansas Academy of Science*, 106: 171-176.
- Shimada K., Everhart M.J., Decker R. & Decker P. D. (2010). A new skeletal remain of the durophagous shark, *Ptychodus mortoni*, from the Upper Cretaceous of North America: an indication of gigantic body size. *Cretaceous Research*, 31: 249-254.
- Shimada K. & Fielitz C. (2006). Annotated checklist of fossil fishes from the Smoky Hill Chalk of the Niobrara Chalk (Upper Cretaceous) in Kansas. In Lucas S.G. & Sullivan R.M. (eds), Late Cretaceous vertebrates from the Western Interior. *New Mexico Museum of Natural History and Science Bulletin*, 35: 193-213.
- Shimada K., Rigsby C.K. & Kim S.H. (2009). Partial skull of Late Cretaceous durophagous shark, *Ptychodus occidentalis* (Elasmobranchii: Ptychodontidae), from Nebraska, USA. *Journal of Vertebrate Paleontology*, 29: 336-349.
- Sirna G., Dalla Vecchia F.M., Muscio G. & Piccoli G. (1994). Catalogue of Paleozoic and Mesozoic vertebrates and vertebrate localities of the Tre Venezie area (North Eastern Italy). *Memorie di Scienze Geologiche, Dipartimento di Geoscienze (Università degli Studi di Padova)*, 46: 255-281.
- Siverson M. (1999). A new large lamniform shark from the uppermost Gearle Siltstone (Cenomanian, Late Cretaceous) of Western Australia. *Transactions of the Royal Society of Edinburgh: Earth Sciences*, 90: 49-66.
- Sorbini L. (1976). L'ittiofauna cretacea di Cinto Euganeo (Padova-Nord Italia). *Bollettino del Museo Civico di Storia Naturale di Verona*, 3: 479-567.
- Sorbini L. (1980). Il giacimento di Cinto Euganeo (Padova). In I vertebrati fossili italiani, Catalogo della Mostra: 119-122.
- Trbušek J. (1999). Upper Cretaceous sharks and rays from the Prokop opencast mine at Březina near Moravská Třebová. *Acta Universitatis Palackianae Olomucensis, Facultas Rerum Naturalium, Geologica*, 36: 51-61.
- Trif N. & Codrea V.A. (2017). Ocurențe ale genului mezozoic *Ptychodus* (Elasmobranchii) în România: date preliminare. *Al XV - lea Simpozion Național Studențesc „GEOECOLOGIA” 2017*: 7-11.

- Underwood C.J. & Cumbaa S.L. (2010). Chondrichthyans from a Cenomanian (Late Cretaceous) bonebed, Saskatchewan, Canada. *Palaeontology*, 53, 903-944.
- Uyeno T. (1972). On Cretaceous and Tertiary fish remain from the Hidaka and Yūbari Districts in Hokkaido, Japan. *Memoirs of the National Science Museum*, 5: 223-226.
- Uyeno T. & Sakura H. (1990). Catalogue of fish fossil specimens. *National Science Museum*. vi + 70 pp. Tokyo.
- Verma O. (2015). Cretaceous vertebrate fauna of the Cauvery Basin, southern India: Palaeodiversity and palaeobiogeographic implications. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 431: 53-67.
- Verma O., Prasad G.V., Goswami A. & Parmar V. (2012). *Ptychodus decurrens* Agassiz (Elasmobranchii: Ptychodontidae) from the Upper Cretaceous of India. *Cretaceous Research*, 33: 183-188.
- Vullo R. & Arnaud E. (2009). Présence de *Ptychodus latissimus* Agassiz, 1843 (Elasmobranchii, Hybodontiformes) dans le Crétacé supérieur des Charentes. *Annales de la Société des Sciences Naturelles de la Charente-Maritime*, 9: 962-966.
- Welton B.J. & Farish R.F. (1993). The collector's guide to fossil sharks and rays from the Cretaceous of Texas. 204 pp. Horton Printing Co., Dallas.
- Wenz S. (1972). Présence du sélacien *Ptychodus* (*Pt. chappelli*) dans le Crétacé supérieur de l'Équateur (Amérique du Sud). *Bulletin du Muséum National d'Histoire Naturelle, Paris, Série 3, Sciences de la Terre*, 13: 91-94.
- Whitham F. (1991). The stratigraphy of the Upper Cretaceous Ferriby, Welton and Burnham formations north of the Humber, north-east England. *Proceedings of the Yorkshire Geological Society*, 48: 227-254.
- Wiese F., Čech S., Ekrť B., Košťák M., Mazuch M. & Voigt S. (2004). The Upper Turonian of the Bohemian Cretaceous Basin (Czech Republic) exemplified by the Úpohlavy working quarry: integrated stratigraphy and palaeoceanography of a gateway to the Tethys. *Cretaceous Research*, 25: 329-352.
- Williamson T.E., Kirkland J.I. & Lucas S.G. (1993). Selachians from the Greenhorn cyclothem („Middle“ Cretaceous: Cenomanian–Turonian), Black Mesa, Arizona, and the paleogeographic distribution of Late Cretaceous selachians. *Journal of Paleontology*, 67: 447-474.
- Wittler F. (1995). Fische aus dem Turon des Dortmunder Stadtgebietes. *Arbeitskreis Paläontologie Hannover*, 23: 41-59.
- Yabumoto Y. & Uyeno T. (1994). Late Mesozoic and Cenozoic fish faunas of Japan. *The Island Arc*, 3: 255-269.
- Zorzin R. (2001). Fossili. Per conoscere il nostro passato. 128 pp. Demetra, Verona.
- Zorzin R. (ed.) (2017). Vertebrati fossili, marini e terrestri del Veronese. 174 pp. Cierre edizioni, Verona.